



congenital adrenal hyperplasia due to 11-beta-hydroxylase deficiency

Congenital adrenal hyperplasia (CAH) due to 11-beta-hydroxylase deficiency is one of a group of disorders (collectively called congenital adrenal hyperplasia) that affect the adrenal glands. The adrenal glands are located on top of the kidneys and produce a variety of hormones that regulate many essential functions in the body. In people with CAH due to 11-beta-hydroxylase deficiency, the adrenal glands produce excess androgens, which are male sex hormones.

There are two types of CAH due to 11-beta-hydroxylase deficiency, the classic form and the non-classic form. The classic form is the more severe of the two types.

Females with the classic form of CAH due to 11-beta-hydroxylase deficiency have external genitalia that do not look clearly male or female (ambiguous genitalia). However, the internal reproductive organs develop normally. Males and females with the classic form of this condition have early development of their secondary sexual characteristics such as growth of facial and pubic hair, deepening of the voice, appearance of acne, and onset of a growth spurt. The early growth spurt can prevent growth later in adolescence and lead to short stature in adulthood. In addition, approximately two-thirds of individuals with the classic form of CAH due to 11-beta-hydroxylase deficiency have high blood pressure (hypertension). Hypertension typically develops within the first year of life.

Females with the non-classic form of CAH due to 11-beta-hydroxylase deficiency have normal female genitalia. As affected females get older, they may develop excessive body hair growth (hirsutism) and irregular menstruation. Males with the non-classic form of this condition do not typically have any signs or symptoms except for short stature. Hypertension is not a feature of the non-classic form of CAH due to 11-beta-hydroxylase deficiency.

Frequency

CAH due to 11-beta-hydroxylase deficiency accounts for 5 to 8 percent of all cases of congenital adrenal hyperplasia. It is estimated that CAH due to 11-beta-hydroxylase deficiency occurs in 1 in 100,000 to 200,000 newborns. This condition is more common in Moroccan Jews living in Israel, occurring in approximately 1 in 5,000 to 7,000 newborns. The classic form of CAH due to 11-beta-hydroxylase deficiency appears to be much more common than the non-classic form.

Genetic Changes

Mutations in the *CYP11B1* gene cause CAH due to 11-beta-hydroxylase deficiency. The *CYP11B1* gene provides instructions for making an enzyme called 11-beta-hydroxylase. This enzyme is found in the adrenal glands, where it helps produce hormones called cortisol and corticosterone. Cortisol has numerous functions, such as maintaining blood sugar levels, protecting the body from stress, and suppressing inflammation. Corticosterone gets converted to the hormone aldosterone, which helps control blood pressure by maintaining proper salt and fluid levels in the body.

CAH due to 11-beta-hydroxylase deficiency is caused by a shortage (deficiency) of the 11-beta-hydroxylase enzyme. When 11-beta-hydroxylase is lacking, precursors that are used to form cortisol and corticosterone build up in the adrenal glands and are converted to androgens. The excess production of androgens leads to abnormalities of sexual development, particularly in females with CAH due to 11-beta-hydroxylase deficiency. A buildup in the precursors used to form corticosterone increases salt retention, leading to hypertension in individuals with the classic form of CAH due to 11-beta-hydroxylase deficiency.

The amount of functional 11-beta-hydroxylase enzyme that an individual produces typically determines the extent of abnormal sexual development. Individuals with the classic form of the condition usually have *CYP11B1* gene mutations that result in the production of an enzyme with low levels of function or no function at all. Individuals with the non-classic form of the condition typically have *CYP11B1* gene mutations that lead to the production of an enzyme with moderately reduced function. The severity of the signs and symptoms of sexual development do not appear to be related to the severity of the hypertension.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- 11 beta hydroxylase deficiency
- 11b hydroxylase deficiency
- adrenal hyperplasia, hypertensive form
- deficiency of steroid 11-beta-monooxygenase
- P450C11B1 deficiency
- steroid 11 beta hydroxylase deficiency

Diagnosis & Management

These resources address the diagnosis or management of congenital adrenal hyperplasia due to 11-beta-hydroxylase deficiency:

- Genetic Testing Registry: Deficiency of steroid 11-beta-monooxygenase
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268292/>
- MedlinePlus Encyclopedia: Congenital Adrenal Hyperplasia
<https://medlineplus.gov/ency/article/000411.htm>

These resources from MedlinePlus offer information about the diagnosis and management of various health conditions:

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Congenital Adrenal Hyperplasia
<https://medlineplus.gov/ency/article/000411.htm>
- Health Topic: Adrenal Gland Disorders
<https://medlineplus.gov/adrenalglanddisorders.html>
- Health Topic: Newborn Screening
<https://medlineplus.gov/newbornscreening.html>

Genetic and Rare Diseases Information Center

- 11-beta-hydroxylase deficiency
<https://rarediseases.info.nih.gov/diseases/5658/11-beta-hydroxylase-deficiency>

Additional NIH Resources

- Eunice Kennedy Shriver National Institute of Child Health and Human Development
<https://www.nichd.nih.gov/health/topics/cah/conditioninfo/Pages/default.aspx>
- National Heart Lung and Blood Institute: What Is High Blood Pressure?
<https://www.nhlbi.nih.gov/health/health-topics/topics/hbp/>
- National Institutes of Health Clinical Center Patient Education
https://www.cc.nih.gov/ccc/patient_education/pepubs/cah.pdf

Educational Resources

- Boston Children's Hospital: Hypertension in Children
<http://www.childrenshospital.org/conditions-and-treatments/conditions/hypertension>
- Centers for Disease Control and Prevention: High Blood Pressure
<https://www.cdc.gov/bloodpressure/>
- Cleveland Clinic: Hypertension
<http://my.clevelandclinic.org/health/articles/hypertension-high-blood-pressure>
- Cleveland Clinic: Sexual Hormone Disorders
<http://my.clevelandclinic.org/health/articles/sexual-hormone-disorders>
- Disease InfoSearch: 11-Beta-Hydroxylase Deficiency
<http://www.diseaseinfosearch.org/11-Beta-Hydroxylase+Deficiency/2>
- Intersex Society of North America
<http://www.isna.org/faq/conditions/cah>
- MalaCards: adrenal hyperplasia, congenital, due to 11-beta-hydroxylase deficiency
http://www.malacards.org/card/adrenal_hyperplasia_congenital_due_to_11_beta_hydroxylase_deficiency
- March of Dimes: Genital and Urinary Tract Defects
<http://www.marchofdimes.org/baby/genital-and-urinary-tract-defects.aspx>
- Merck Manual Consumer Version: The Body's Control of Blood Pressure
<http://www.merckmanuals.com/home/heart-and-blood-vessel-disorders/high-blood-pressure/high-blood-pressure>
- Merck Manual Home Edition for Patients and Caregivers: Virilization
<http://www.merckmanuals.com/home/hormonal-and-metabolic-disorders/adrenal-gland-disorders/virilization>

- Orphanet: Congenital adrenal hyperplasia due to 11-beta-hydroxylase deficiency
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=90795
- Virginia Department of Health
http://www.vdh.virginia.gov/content/uploads/sites/33/2016/11/Parent-Fact-Sheet_CAH_English.pdf

Patient Support and Advocacy Resources

- American Heart Association: High Blood Pressure
http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/High-Blood-Pressure-ATH_UCM_002020_SubHomePage.jsp
- CARES Foundation
<http://www.caresfoundation.org/>
- CLIMB: Children Living with Inherited Metabolic Diseases
<http://www.climb.org.uk/>
- National Organization for Rare Disorders (NORD)
<https://rarediseases.org/rare-diseases/congenital-adrenal-hyperplasia/>
- The MAGIC Foundation
<https://www.magicfoundation.org/Growth-Disorders/Congenital-Adrenal-Hyperplasia/>

Genetic Testing Registry

- Deficiency of steroid 11-beta-monooxygenase
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268292/>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22congenital+adrenal+hyperplasia+due+to+11-beta-hydroxylase+deficiency%22+OR+%22Congenital+Adrenal+Hyperplasia%22+OR+%22Adrenal+Hyperplasia%2C+Congenital%22>

Scientific articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Adrenal+Hyperplasia,+Congenital%5BMAJR%5D%29+AND+%28%2811+beta+hydroxylase+deficiency%5BTIAB%5D%29+OR+%28steroid+11+beta+hydroxylase+deficiency%5BTIAB%5D%29+OR+%2811+hydroxylase+deficiency%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- ADRENAL HYPERPLASIA, CONGENITAL, DUE TO STEROID 11-BETA-HYDROXYLASE DEFICIENCY
<http://omim.org/entry/202010>

Sources for This Summary

- Krone N, Riepe FG, Götze D, Korsch E, Rister M, Commentz J, Partsch CJ, Grötzinger J, Peter M, Sippell WG. Congenital adrenal hyperplasia due to 11-hydroxylase deficiency: functional characterization of two novel point mutations and a three-base pair deletion in the CYP11B1 gene. *J Clin Endocrinol Metab*. 2005 Jun;90(6):3724-30. Epub 2005 Mar 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15755848>
- Nimkarn S, New MI. Steroid 11beta- hydroxylase deficiency congenital adrenal hyperplasia. *Trends Endocrinol Metab*. 2008 Apr;19(3):96-9. doi: 10.1016/j.tem.2008.01.002. Epub 2008 Feb 21. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18294861>
- Parajes S, Loidi L, Reisch N, Dhir V, Rose IT, Hampel R, Quinkler M, Conway GS, Castro-Feijóo L, Araujo-Vilar D, Pombo M, Dominguez F, Williams EL, Cole TR, Kirk JM, Kaminsky E, Rumsby G, Arlt W, Krone N. Functional consequences of seven novel mutations in the CYP11B1 gene: four mutations associated with nonclassic and three mutations causing classic 11{beta}-hydroxylase deficiency. *J Clin Endocrinol Metab*. 2010 Feb;95(2):779-88. doi: 10.1210/jc.2009-0651. Epub 2010 Jan 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20089618>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2846960/>
- Peter M. Congenital adrenal hyperplasia: 11beta-hydroxylase deficiency. *Semin Reprod Med*. 2002 Aug;20(3):249-54. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12428205>
- Zhu YS, Cordero JJ, Can S, Cai LQ, You X, Herrera C, DeFillo-Ricart M, Shackleton C, Imperato-McGinley J. Mutations in CYP11B1 gene: phenotype-genotype correlations. *Am J Med Genet A*. 2003 Oct 15;122A(3):193-200.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12966519>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/congenital-adrenal-hyperplasia-due-to-11-beta-hydroxylase-deficiency>

Reviewed: March 2011

Published: January 24, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services